
G٥



☑e-mail 🚇 print

My Profile Log in

InterScience\*

Home / Polymers and Materials Science / Polymers and Materials Science (general)



Journal of Polymer Science

Volume 4, Issue 6 , Pages 673 - 688

Published Online: 10 Mar 2003

Copyright @ 1949 Interscience Publishers, Inc.

Save Title to My Profile

Set E-Mall Alert

So to the homepode for this journal to access triels, semple copies, ed total and author

information, news, and more,

IN THIS TITLE

All Content

Publication Titles

Journal of Polymer Science

CrossRef / Google Search

Advanced Search

Acronym Finder

All Fields

PHOME 4

PAROUT US

**Download Citation** < Previous Abstract | Next Abstract > Abstract | References | Full Text: PDF (997k) | Related Articles | Citation Tracking

## Article

Water content of hydrophilic phenoi-formaldehyde resins: Vapor pressure-temperature relationships

Ralph W. Bentz, Harvey A. Neville

Save Article to My Profile

Department of Chemistry, Lehigh University, Bethlehem, Pennsylvania

## **Abstract**

The significance of the water content to the physical properties of phenol-formaldehyde resins at various stages of preparation is briefly discussed. Available methods for the determination of water in such resins were found to be either unrellable or too elaborate for process control. It was undertaken to develop a simple, rapid, relative method for the continuous measurement of water content which can be used to follow the follow the process of dehydration and to indicate the end-point which will result in the best dispersion of water in the cured resin. In order to standardize or calibrate such a relative method, it was necessary also to develop an absolute method for the analysis of water in the resol.

The composition of a homogeneous mixture of liquids is reflected by its vapor pressuretemperature relationships. During dehydration at 80°C., the vapor pressure of the resol decreases about 350 mm. of mercury. A very sharp increase in the slope of the vapor pressure curve occurs at approximately the proper degree of dehydration. At this stage a slight decrease in water content causes a very large decrease in vapor pressure. The sharp inflection in the curve is attributed to the existence of both "free" and "bound" water in the resin.

During the process of vacuum dehydration of the resin, while the "free" water decreases to approximately one-eighth of its original content, the "bound" water remains substantially constant. The binding of water is attributed to the methylol groups of the resol molecules and its release during the cure or hardening of the resin is considered to result from the conversion of these groups to methylene linkages.

Received: 24 January 1949 Digital Object Identifier (DOI)

10.1002/pol.1949.120040601 About DQI

## Related Articles

- Find other articles like this in Wiley InterScience
- Find articles in Wiley InterScience written by any of the authors

BY CITATION

Vol: | Go |

Journal of Polymer Science, Part B: Polymer Physics 2006 Prize Winners

V. Kuppa, S. Menakanit, R. Krishnamoorti, E. Manias for their paper?

?Simulation Insights on the Structure of Nanoscopically Confined Poly(ethylene oxide)?

Read the article

Journal of Polymer Science, Part A: Polymer Chemistry

Impact Factor increased to

2.773

One of the top Journals in Polymer Science!

\* ISI Journal Citation Report 2004

Wiley InterScience is a member of CrossRef.



Go to journal



Impact Factor increased to

2.714 #23 among 125 journals in the category of Chemistry. Multidisciplinary

\* ISI Journal Citation Report 2004 Go to journal

HIGHLIGHT **ARTICLES** Journal of Polymer Science, Part A: Polymer Chemistry Highlight Articles are specially commissioned, providing a unique insight into the research activities and findings of the top laboratories throughout the world.

Free online access

About Wiley InterScience | About Wiley | Privacy | Terms & Conditions Copyright @ 1999-2006 John Wiley & Sons, Inc. All Rights Reserved.